

1 Type or print clearly:

2 NAME: Bill Burgos

3 AGENCY/COMPANY: Penn State

4 ADDRESS: Department of Civil and Environmental Engineering, 212 Sackett Bldg

5 CITY, STATE, ZIP: University Park, PA 16802

6 TELEPHONE: 814-836-0578 FAX: 814-863-7304

7 E-MAIL : wdb3@psu.edu

8 Type of presentation: Oral Poster

9 **Technical Division or Session:**

10 Ecology Forestry and Wildlife Geotechnical Engineering

11 Refuse and Tailings Management & Reclamation Land Use and Design

12 Soils and Overburden Water Management Other Case Study

13 Other interests for those persons who may or may not be presenting a paper:

14 Field Tour Workshop Send Hotel and Conference Registration information

15

16 Exhibitor Attendee

17 Sponsor (Gold, Silver, Bronze or Break)

18

19 **Importance of fungi in biological Mn(II) oxidation in limestone treatment beds**

20 William D. Burgos¹, Hui Tan¹, Cara M. Santelli² and Colleen M. Hansel²

21 ¹Department of Civil and Environmental Engineering, The Pennsylvania State University,

22 University Park, PA, USA. ²Harvard School of Engineering and Applied Sciences, Cambridge,

23 **ABSTRACT**

24 Coal mine drainage (CMD) is the single greatest threat to the environment of Appalachia
25 in the United States, releasing large volumes of acid and metal contaminants (e.g., Mn, Fe).

26 Passive limestone-based treatment systems are the most cost-effective method for manganese(II)
27 removal from CMD. The success of passive Mn(II)-removal systems has been variable due to a
28 poor understanding of the mechanisms of Mn removal and the microbial communities involved.

29 We selected one Mn-removal system from northwestern Pennsylvania that treats an

30 exceptionally high concentration of 150 mg/L Mn(II), and conducted laboratory experiments to
31 evaluate the relative importance of abiotic versus biotic processes responsible for Mn removal,
32 and to evaluate the relative importance of bacteria versus fungi on biological Mn(II) oxidation.

33 We found that while abiotic processes such as Mn(II) sorption and heterogeneous oxidation
34 contribute to Mn removal, biological Mn(II) oxidation is the most important process to ensure
35 effective, long-term Mn removal. We also found, that fungal activity accounted for over 80 % of
36 Mn(II) oxidation in this Mn-removal bed. We also selected four additional Mn-removal systems
37 from western Pennsylvania for an extensive culture enrichment survey of bacteria and fungi.

38 From this survey, we found that Mn(II)-oxidizing fungi were isolated more readily than Mn(II)-
39 oxidizing bacteria – fungal isolates outnumbered bacterial isolates 84:10 in 3 of the 4 systems,
40 and that fungi were extremely tolerant to elevated concentrations of Mn(II).

41 **Key Words:** acid mine drainage, metal oxidation, manganese removal

42 INTRODUCTION

43 The removal of elevated concentrations of dissolved manganese, Mn(II), from mine
44 drainage is a significant problem for many regions in the United States and throughout the world.
45 In Appalachia, centuries of coal mining have left thousands of abandoned mines that are
46 discharging metal-contaminated effluent, with Mn concentrations as high as 150 mg/L, and
47 severely degrading water resources (Cravotta III, 2007 and references therein; Herlihy et al.,
48 1990). Manganese, while not considered to be acutely toxic to humans (although this is poorly
49 studied), can be damaging to ecosystems and water distribution networks. Furthermore, Mn(II)
50 oxidation reactions produce protons, which perpetuates the generation of acid mine drainage
51 (AMD). In the eastern United States, one of the most common methods to remediate high
52 concentrations of dissolved Mn(II) in coal mine drainage (CMD) is the use of biologically-active
53 limestone treatment beds, where oxidation and subsequent precipitation of Mn(III/IV) oxides is
54 catalyzed by abiotic and biotic processes (Hallberg and Johnson, 2005; Johnson et al., 2005). In
55 essence, Mn is immobilized via the precipitation of sparingly soluble minerals, which
56 subsequently remove other metal contaminants (e.g., Cu, Co, Zn) through both co-precipitation
57 and surface adsorption reactions. To date, the overall success of Mn removal within passive
58 treatment systems is widely variable due to a poor understanding of the processes and
59 mechanisms that govern Mn(II) oxidation at near-neutral pH in these systems.

60 Microorganisms accelerate Mn(II) oxidation rates up to several orders of magnitude
61 faster than abiotic catalysis, thus it is believed that the precipitation of Mn(III/IV) oxide minerals
62 in the environment is largely driven by microbiological activity (Nealson et al., 1988; Tebo,
63 1991; Tebo et al., 2004). Mn(II)-oxidizing bacteria are ubiquitously distributed in the
64 environment and much research has recently been devoted to understanding the mechanisms,

65 pathways, and products of Mn(II) oxidation by bacteria (Bargar et al., 2005; Dick et al., 2008b;
66 Francis et al., 2001; Hansel and Francis, 2006; Johnson and Tebo, 2007; Ridge et al., 2007;
67 Webb et al., 2005a; Webb et al., 2005b). Although much less studied, Mn(II)-oxidizing fungi
68 have also been recovered from a wide variety of ecosystems, such as agricultural soil (Pedler et
69 al., 1996), deep sea sediments (Shao and Sun, 2007), building stone, (de la Torre and Gomez-
70 Alarcon, 1994), desert varnish (Grote and Krumbein, 1992; Krumbein and Jens, 1981),
71 streambeds (Miyata et al., 2006a; Miyata et al., 2004; Takano et al., 2006), and an artificial
72 wetland (Takano et al., 2006).

73 The contribution of microbial activity in the remediation of Mn-contaminated waters has
74 frequently been observed (Bamforth et al., 2006; Haack and Warren, 2003; Hallberg and
75 Johnson, 2005; Johnson et al., 2005; Johnson and Younger, 2005). Several different strains of
76 Mn(II)-oxidizing bacteria have even been used in a patented bioremediation method, the
77 “Pyrolusite Process”, for treating manganiferous mine waters (Vail and Riley, 2000). A recent
78 study by Mariner et al. (2008) identified Mn(II)-oxidizing fungi, in addition to bacteria, that
79 successfully grow in a Mn(II) attenuation bioreactor for treating mine waters. To our
80 knowledge, this is the first study to document the role of fungi in these bioremediation
81 technologies. In general, however, the identities and growth characteristics of the Mn(II)-
82 oxidizing community contributing to Mn remediation remains largely unresolved.

83 The purpose of this research was to improve our understanding and design capability of
84 passive manganese(II)-removal systems for the treatment of coal mine drainage. The objectives
85 of this research were to: 1) Measure Mn(II)-removal kinetics in controlled laboratory
86 experiments; and 2) Characterize the microbial communities that promote Mn(II) oxidation in
87 these Mn-removal systems using a culture-based approach.

88 **MATERIALS AND METHODS**

89 **Site description and sampling techniques**

90 One passive Mn-removal system, referred to as Fairview, was characterized in this study.
91 The system is located in northwestern Pennsylvania (Elk County, 41°21'05" N, 78°39'16" W)
92 and was constructed in 2004 to treat CMD from a surface coal mine with an influent Mn(II)
93 concentration of 130 – 150 mg/L. CMD is conveyed through an underground limestone drain
94 before discharging into a limestone-laden constructed wetland and then enters the limestone Mn-
95 removal bed. Discharge into the wetland comes from a pipe elevated several feet above the
96 ground that provides aeration as the water enters the wetland. Iron is completely removed before
97 entering the Mn-removal bed (Table 1). The rectangular bed (10-cm limestone gravel, 30-m
98 long, 15-m wide, 1-m deep) contains five ditches (perpendicular to bed flow, ~½ the depth of the
99 bed) that were added to promote passive aeration and served as our water and sediment sampling
100 locations. Straw and corncobs were worked into the Mn-removal bed after initial construction
101 when the ditches were added. The flow rate into the Mn-removal bed is ca. 10 gallons/min and
102 the bed provides a hydraulic residence time of ca. 100 hrs.

103 Water and sediment samples were collected in July, September, and November of 2005,
104 April, June, September and December of 2006, and May of 2007 to monitor system performance
105 during all seasons and over an extended period of time. Samples were collected from the
106 influent and effluent pipes, the wetland and from the five ditches in the Mn-removal bed. Water
107 samples were filtered (0.2 µm) in the field and chemically preserved (dependent on analyte).
108 Sediment “crust” samples were collected with sterile spatulas and placed in sterile endotoxin-free
109 centrifuge tubes or whirl-pak bags. Sediments were scraped from the top 1-cm of black
110 precipitates that armored the limestone cobbles. All samples were stored on ice for transport to

111 the laboratory, except for samples intended for nucleic acid-based microbial community analysis,
112 which were transported on dry ice. Dissolved oxygen (DO) concentrations, temperature, pH and
113 conductivity were determined in the field using portable meters.

114 **Sediment incubation experiments**

115 Laboratory sediment incubation experiments were conducted to determine the relative
116 importance of biological activity on Mn(II) loss from solution. Experiments were conducted
117 with a single, large quantity of sediment collected from the middle ditches in the Mn-removal
118 bed. Sediment crusts were collected with sterile spatulas, homogenized, and wet sieved (<2-mm
119 sieve fraction), using sterile tools in the field, into sterile mason jars. Experiments were
120 conducted with 1.0 g moist “live” sediments (0.60 g dry mass) mixed with 50 mL of filter
121 sterilized influent water in 120 mL serum bottles with air or 100 % N₂ in the headspace and
122 sealed with thin Teflon-coated stoppers and Al crimp tops. Abiotic controls were conducted
123 with “killed” sediments that been exposed to 100 kilogray of ⁶⁰Co γ irradiation and maintained
124 under a 100 % N₂ headspace. No loss of Mn(II) was measured in sediment-free controls under
125 an air headspace. All reactors were kept in the dark and shaken at 100 rpm at room temperature
126 (25 °C). Water and sediment suspension samples were periodically collected from the reactors
127 using sterile needle and syringe to measure soluble Mn(II) and pH. Samples were centrifuged at
128 11,000 g for 10 minutes to separate liquid and sediments. Mn(II) concentration in the
129 supernatants were measured using the PAN method (Goto et al., 1976). pH was measured using
130 a combination electrode (Thermo Scientific, Waltham, MA). When the soluble Mn(II)
131 concentration decreased to below detection limit (<0.1 mg/L) in the “live” reactors, MnCl₂ was
132 re-spiked into the reactors to re-establish a Mn(II) concentration of ca. 150 mg/L and MnCl₂ was
133 spiked into the reactors three times.

134 **Culture enrichments and isolations**

135 Samples for culture enrichments were collected in October 2007 from four different
136 treatment systems near Fairview: Saxman Run (Westmoreland County, PA), De Sale Phase 1, De
137 Sale Phase II, and De Sale Phase III (Butler County, PA) as described previously (Santelli et al.,
138 2010). Rock and sediment samples were lightly crushed and homogenized using a mortar and
139 pestle. All samples were diluted in sterile, artificial freshwater (AFW; 34.22 mM NaCl, 1.62
140 mM MgSO₄*7H₂O, 1.36 mM CaCl₂*2H₂O, 0.44 mM K₂HPO₄*3H₂O, 20 mM HEPES buffer at
141 pH7) using serial dilutions to 1/10⁴. Dilutions were plated onto 7 different types of agar-
142 solidified media with 20 mM HEPES buffer (pH 7) and 200 μM MnCl₂. The culture media were
143 described previously: HEPES-buffered AY medium (Miyata et al., 2004); K, Leptothrix, and M
144 media with 0.2 μm filter-sterilized natural site water (Templeton et al., 2005); J and J+acetate
145 (JAC) media with AFW (Hansel and Francis, 2006); and Medium 3 (M3; de la Torre and
146 Gomez-Alarcon, 1994). Mn(II)-oxidizing bacteria and fungi were putatively identified by the
147 presence of brown/black precipitates, Mn(III/IV) oxides, and transferred to fresh media a
148 minimum of 5 times until cultures were deemed axenic. Mn(III/IV) was confirmed using the
149 leucoberbelin blue (LBB) colorimetric assay – LBB reacts specifically with Mn(III) and Mn(IV)
150 and turns deep blue (Krumbein and Altman, 1973).

151 **Growth experiments**

152 Fungal isolates were tested for their ability to grow and oxidize Mn(II) in the presence of
153 varying metal concentrations. Fungi were grown in dark conditions on AY media supplemented
154 with Mn²⁺ (added as MnCl₂) at the following concentrations: 0 μM, 250 μM, 500 μM, 750 μM, 1
155 mM, 5 mM, and 10 mM. Fungi were inoculated with a toothpick in the center of petri dishes and
156 allowed to grow radially outward until the diameter of mycelia reached the plate edges (typically

157 less than 10 days). Colony growth diameter was evaluated approximately every other day. The
158 effects of light/dark and pH (5.5, 6.0, 6.5, 7.0, 7.6, and 8.0) on growth were also tested using the
159 same protocol. MES buffer (20 mM) was used to maintain a solution pH of 5.5, 6.0, and 6.5, and
160 HEPES buffer (20 mM) was used for pH values ≥ 7.0 .

161 **RESULTS AND DISCUSSION**

162 **Mn(II) removal in the field**

163 Limestone beds designed for Mn removal from CMD create a unique geochemical and
164 physical environment for promoting biological Mn(II) oxidation. To function most effectively,
165 Fe, Al and some of the influent acidity are removed first through limestone-based systems before
166 entering the Mn-removal system. In a survey of 140 abandoned coal mine discharges in
167 Pennsylvania, the median Mn concentration was 2.35 mg/L with a range from 0.019 to 74.0
168 mg/L (Cravotta, 2008). Most of these discharges were from underground mines. For the surface
169 coal mine at the Fairview site, the exceptionally high Mn concentration (130 – 150 mg/L) is
170 believed to originate from reactions with the overburden (e.g., MnCO_3 inclusions in limestone).
171 Influent CMD to the Fairview Mn-removal system contains essentially no Fe, Ca and Mg are
172 elevated from upstream neutralization with limestone, and the elevated concentrations of Mn and
173 Al make the water net acidic (Table 1). Sulfate is the primary anion and nutrient concentrations
174 (i.e. TOC, N, P) are relatively low. We have sampled this system on several occasions between
175 December 2005 and May 2007 to capture seasonal effects and have found that Mn is never
176 completely removed through the limestone bed (Tan et al., 2010). Greater removal of Mn(II)
177 occurs in summer months (e.g., 35 mg/L effluent Mn), as compared to the winter sampling event
178 (e.g., 70 mg/L).

179

180 **Mn(II) removal via biotic versus abiotic processes**

181 Reactors containing wet-sieved (<2-mm), MnO_x-rich sediment crusts and their associated
182 natural microbial communities were operated in a fed-batch mode with respect to dissolved
183 Mn(II). MnCl₂ was periodically re-spiked into the reactors to re-establish the initial, influent
184 Mn(II) concentration of ca. 2.5 mM. This fed-batch approach was required to overcome abiotic
185 Mn(II) removal processes such as non-oxidative sorption and heterogeneous oxidation so that the
186 contribution of biological Mn(II) oxidation on Mn removal could be better ascertained. Control
187 reactors that had been sterilized by ⁶⁰Co γ-irradiation and/or maintained under a N₂ atmosphere
188 were compared to “live” (i.e. non-sterile) reactors. Our simplified, operational interpretations
189 were that Mn(II) could be removed only by non-reductive sorption in sterilized reactors
190 maintained under N₂, while both heterogeneous oxidative precipitation of Mn(II) (as MnO_x) and
191 non-reductive sorption could occur in sterilized reactors maintained under air.

192 As seen with the live reactors maintained under air or under N₂ (black and white squares,
193 respectively, in Figure 1), the difference between Mn(II) removal through each fed-batch cycle
194 increased as the experiment proceeded. Our interpretation of these results is that the non-
195 reductive Mn(II) sorption capacity of the sediments was essentially saturated after four cycles.
196 The growing difference between the live reactors maintained under air versus the live reactors
197 under N₂ was presumably caused by biological Mn(II) oxidation and the subsequent precipitation
198 of MnO_x.

199 A series of similar, fed-batch experiments were conducted to determine the effects of
200 dissolved oxygen on Mn(II) removal (Figure 2). A significant difference in Mn(II) removal was
201 observed when the headspace P_{O2} was maintained at 1 % (v/v)-as compared to 10 and 21 %. The
202 difference became more pronounced after several cycles as the abiotic removal mechanisms

203 contributed less to Mn(II) removal (due to saturation). The rate of biological Mn(II) oxidation
204 by *Leptothrix discophora* SS1 has been shown to be directly proportional to dissolved oxygen
205 (Zhang et al., 2002). Our results suggest a different response for the whole, natural microbial
206 community in this Mn-removal bed as compared to *Leptothrix*. Specifically, our results suggest
207 that dissolved oxygen had more of a saturation-type effect on Mn(II) removal where the rate and
208 extent of Mn(II) removal were not different above a P_{O₂} level of 10 %.

209 Fed-batch experiments were also conducted to determine the effects of organic carbon
210 addition on Mn(II) removal (Figure 3). In practice, Mn-removal beds are often designed with an
211 upstream wetland. Dissolved organic carbon (DOC) released via primary production in the
212 wetland into the Mn-removal bed presumably will increase microbial Mn(II) oxidation. Glucose
213 and carboxymethylcellulose (CMC) were selected to represent labile and recalcitrant DOC,
214 respectively. CMC was also selected to represent DOC that might be generated from wood
215 dissolution with the thought that wetlands could possibly be replaced with a solid-phase carbon
216 source emplaced within the bed. The addition of glucose was shown to slightly improve Mn(II)
217 removal while the addition of CMC was shown to significantly inhibit Mn(II) removal. Our
218 interpretation is that glucose stimulated general heterotrophic activity that in turn fortuitously
219 stimulated biological Mn(II) oxidation. We are unsure about the inhibitory effect of CMC,
220 however, we speculate that CMC may complex Mn(II) to form soluble species that are difficult
221 to oxidize.

222 **Mn(II) oxidation via bacterial versus fungal activity**

223 A series of fed-batch experiments were conducted in the presence/absence of fungicides
224 (0.2 g/L cyclohexamide plus 0.2 g/L pentachloronitrobenzene) to operationally assess the relative
225 importance of Mn(II) oxidation by fungi as compared to other microbial community members.

226 A somewhat surprising and important finding from these experiments was that the fungicides
227 knocked out the majority of the sediment's ability to remove Mn(II). Reactors that contained
228 sterile sediments and/or were maintained under N₂ all yielded results similar to those reactors
229 that contained fungicides. Our interpretation of these results is that, e.g., the difference between
230 the live reactors with fungicides (black circles in Figure 4) and the live reactors without
231 fungicides (black squares) was due to fungal activity. Furthermore, we believe that fungi were
232 the most important bio-catalysts of Mn(II) oxidation in this treatment system.

233 Based on enumerations of culturable heterotrophic bacteria measured at the end of these
234 experiments, the fungicides did not non-specifically harm the bacterial population in the
235 sediments (Figure 5). We actually observed an increase in culturable bacterial numbers and
236 speculate that this occurred because of the absence of fungi competing for space on the plates.
237 Another possible explanation for increased numbers of culturable bacterial in the presence of
238 fungicides is that fungi themselves may produce antibiotics to suppress bacterial activity.

239 **Mn(II) oxidation by pure cultures of fungi**

240 We conducted an extensive culture enrichment survey of several Mn(II)-treatment
241 systems in western Pennsylvania. An interesting and unexpected result from this survey was that
242 Mn(II)-oxidizing fungi were isolated more readily than Mn(II)-oxidizing bacteria, where fungal
243 isolates outnumbered bacterial isolates 84 to 10. In other words, in these Mn(II)-treatment
244 systems we found that fungi constituted 89 % of the Mn(II)-oxidizing cultures while bacteria
245 constituted just 11 %. The most commonly recovered species, in terms of number of treatment
246 systems and number of times obtained, was representative isolate DS2psM2a2 (Figure 6, left).
247 Based on phylogenetic analysis and morphological characterization, DS2psM2a2 was identified
248 as *Plectosphaerella cucumerina* belonging to the class Sordariomycetes (Santelli et al., 2010).

249 We initially hypothesized that Mn(II) oxidation was predominantly mediated by bacteria;
250 however, our results suggest that fungi are also important mediators of Mn(II) oxidation. Also of
251 great practical importance, we found that most of these fungi were highly tolerant to elevated
252 Mn(II) concentrations. As shown in Figure 6, two representative fungal isolates from these Mn-
253 removal limestone beds can tolerate astonishingly high levels of Mn(II) (e.g., 1,000 to >10,000
254 μM). It is generally believed that fungi are more tolerant than bacteria to high concentrations of
255 heavy metals leading to a prevalence of fungi in heavy metal-contaminated soil (Chander et al.,
256 2001a; Chander et al., 2001b; Kelly et al., 1999; Rajapaksha et al., 2004). Considering that
257 Mn(II) concentrations in surface coal mine drainage in Appalachia often exceed 1,000 μM (= 55
258 mg/L) and can even exceed 3,000 μM , microbial catalysts in these Mn(II)-removal systems must
259 be tolerant of high Mn(II) concentrations – further demonstrating that Mn(II)-oxidizing fungi
260 likely play an important role in the treatment process. Consistent with this conclusion, a recent
261 study (Mariner et al., 2008) suggested that Mn(II)-oxidizing fungi are more abundant than
262 bacteria in bioreactors treating mine water with elevated Mn(II) concentrations.

263

264 **Acknowledgements:**

265 This research was supported by the National Science Foundation through Grant No. CHE-
266 0431328 and Grant No. EAR07-45374, and by the Office of Surface Mining under Cooperative
267 Agreement S07AP12478.

268

269 **References**

- 270 Bamforth S, Manning D, Singleton I, Younger P, Johnson K (2006). Manganese removal from mine
271 waters—investigating the occurrence and importance of manganese carbonates. *Applied*
272 *geochemistry* **21**: 1274-1287.
- 273 Bargar JR, Tebo BM, Bergmann u, Webb SM, Glatzel P, Chiu VQ *et al* (2005). Biotic and abiotic products
274 of Mn(II) oxidation by spores of the marine *Bacillus sp.* strain SG-1. *American Mineralogist* **90**:
275 143-154.
- 276 Chander K, Dyckmans J, Hoepfer H, Joergensen R, Raubuch M (2001a). Long-term effects on soil
277 microbial properties of heavy metals from industrial exhaust deposition. *Journal of Plant*
278 *Nutrition and Soil Science* **164**: 657 - 663.
- 279 Chander K, Dyckmans J, Joergensen R, Meyer B, Raubuch M (2001b). Different sources of heavy metals
280 and their long-term effects on soil microbial properties. *Biology and Fertility of Soils* **34**: 241-247.
- 281 Cravotta, C.A., 2008. Dissolved metals and associated constituents in abandoned coal-mine discharges,
282 Pennsylvania, USA—1. Constituent concentrations and correlations. *Appl. Geochem.* **23**: 203-
283 226.
- 284 de la Torre MA, Gomez-Alarcon G (1994). Manganese and iron oxidation by fungi isolated from building
285 stone. *Microb Ecol* **27**: 177-188.
- 286 Dick GJ, Torpey JW, Beveridge TJ, Tebo BM (2008). Direct Identification of a Bacterial Manganese(II)
287 Oxidase, the Multicopper Oxidase MnxG, from Spores. *Appl Environ Microbiol* **74**: 1527-1534.
- 288 Francis C, Co E, Tebo B (2001). Enzymatic Manganese (II) Oxidation by a Marine α -Proteobacterium.
289 *Applied and Environmental Microbiology* **67**: 4024-4029.
- 290 Goto, K., Taguchi, S., Fukue, Y., Ohta, K., Watanabe, H., 1977. Spectrophotometric determination of
291 manganese with 1-(2-pyridylazo)-2-naphthol and a non-ionic surfactant. *Talanta*. **24**: 752-753.
- 292 Grote G, Krumbein W (1992). Microbial precipitation of manganese by bacteria and fungi from desert
293 rock and rock varnish. *Geomicrobiology Journal* **10**: 49-57.
- 294 Haack EA, Warren L (2003). Biofilm hydrous manganese oxyhydroxides and metal dynamics in acid rock
295 drainage. *Environ Sci Technol* **37**: 4138-4147.
- 296 Hallberg K, Johnson D (2005). Biological manganese removal from acid mine drainage in constructed
297 wetlands and prototype bioreactors. *Science of the Total Environment* **338**: 115-124.
- 298 Hansel CM, Francis CA (2006). Coupled photochemical and enzymatic Mn(II) oxidation pathways of a
299 planktonic *Roseobacter*-like bacterium. *Applied and Environmental Microbiology* **72**: 3543-3549.

300 Herlihy A, Kaufmann P, Mitch M, Brown D (1990). Regional estimates of acid mine drainage impact on
301 streams in the Mid-Atlantic and Southeastern United States. *Water, Air, & Soil Pollution* **50**: 91-
302 107.

303 Johnson H, Tebo B (2007). In vitro studies indicate a quinone is involved in bacterial Mn (II) oxidation.
304 *Arch Microbiol* **189**: 59-69.

305 Johnson K, Baker A, Manning D (2005). Passive treatment of Mn-rich mine water: Using fluorescence to
306 observe microbiological activity. *Geomicrobiology Journal* **22**: 141-149.

307 Johnson KL, Younger PL (2005). Rapid Manganese Removal from Mine Waters Using an Aerated Packed-
308 Bed Bioreactor. *J Environ Qual* **34**: 987-993.

309 Kelly JJ, Hogblom M, Tatellii RL (1999). Changes in soil microbial communities over time resulting from
310 one time application of zinc: a laboratory microcosm study. *Soil Biology and Biochemistry* **31**:
311 1455-1465.

312 Krumbein WE, Altman HJ (1973). A new method for detection and enumeration of manganese-oxidizing
313 and -reducing microorganisms. *Helgol Wiss Meeresunters* **25**: 4985-4999.

314 Krumbein WE, Jens K (1981). Biogenic rock varnishes on the Negev Desert (Israel): an ecological study of
315 iron and manganese transformation by cyanobacteria and fungi. *Oecologia* **50**: 25-38.

316 Mariner R, Johnson D, Hallberg K (2008). Characterisation of an attenuation system for the remediation
317 of Mn (II) contaminated waters. *Hydrometallurgy* **94**: 100-104.

318 Miyata N, Maruo K, Tani Y, Tsuno H, Seyama H, Soma M *et al* (2006). Production of Biogenic Manganese
319 Oxides by Anamorphic Ascomycete Fungi Isolated from Streambed Pebbles. *Geomicrobiology*
320 *Journal* **23**: 63-73.

321 Miyata N, Tani Y, Iwahori K, Soma M (2004). Enzymatic formation of manganese oxides by an
322 Acremonium-like hyphomycete fungus, strain KR 21-2. *FEMS Microbiology Ecology* **47**: 101-109.

323 Nealson K, Tebo B, Rosson R (1988). Occurrence and mechanisms of microbial oxidation of manganese.
324 *Advances in Applied Microbiology* **33**: 279-318.

325 Pedler J, Webb M, Buchhorn S, Graham R (1996). Manganese-oxidizing ability of isolates of the take-all
326 fungus is correlated with virulence. *Biology and Fertility of Soils* **22**: 272-278.

327 Rajapaksha RMCP, Tobor-Kaplon MA, Baath E (2004). Metal Toxicity Affects Fungal and Bacterial
328 Activities in Soil Differently. *Appl Environ Microbiol* **70**: 2966-2973.

329 Ridge J, Lin M, Larsen E, Fegan M, McEwan A, Sly L (2007). A multicopper oxidase is essential for
330 manganese oxidation and laccase-like activity in *Pedomicrobium* sp. ACM 3067. *Environmental*
331 *Microbiology* **9**: 944-953.

332 Santelli, C.M., D.H. Pfister, D. Lazarus, L. Sun, H. Tan, W.D. Burgos, C.M. Hansel. 2010. Characterization
333 of Mn(II)-oxidizing microbial communities in passive remediation systems treating coal mine
334 drainage reveals a role for fungi in metal removal. submitted to *Applied Environmental*
335 *Microbiology* November 2009.

336 Shao Z, Sun F (2007). Intracellular sequestration of manganese and phosphorus in a metal-resistant
337 fungus *Cladosporium cladosporioides* from deep-sea sediment. *Extremophiles* **11**: 435-443.

338 Takano K, Itoh Y, Ogino T, Kurosawa K, Sasaki K (2006). Phylogenetic analysis of manganese-oxidizing
339 fungi isolated from manganese-rich aquatic environments in Hokkaido, Japan. *Limnology* **7**: 219-
340 223.

341 Tan, H., G. Zhang, P.J. Heaney, S.M. Webb and W.D. Burgos. 2010. "Characterization of manganese
342 precipitates from Appalachian coal mine drainage treatment systems." *Applied Geochemistry*, in
343 press 01/2010.

344 Tebo B (1991). Manganese (II) Oxidation in the Suboxic Zone of the Black Sea. *Deep-Sea Research* **38**:
345 S883-S906.

346 Tebo BM, Bargar JR, Clement BG, Dick GJ, Murray KJ, Parker D *et al* (2004). Biogenic manganese oxides:
347 properties and mechanisms of formation. *Annu Rev Earth Planet Sci* **32**: 287-328.

348 Templeton AS, Staudigel H, Tebo BM (2005). Diverse Mn(II)-oxidizing bacteria isolated from submarine
349 basalts at Loihi Seamount. *Geomicrobiology Journal* **22**: 127-139.

350 Vail W, Riley R (2000). The Pyrolusite Process: A bioremediation process for the abatement of acid mine
351 drainage. *Green Lands* **30**: 40-46.

352 Webb S, Dick G, Bargar J, Tebo B (2005a). Evidence for the presence of Mn (III) intermediates in the
353 bacterial oxidation of Mn (II). *Proceedings of the National Academy of Sciences* **102**: 5558-5563.


354 Webb SM, Tebo BM, Bargar JR (2005b). Structural characterization of biogenic Mn oxides produced in
355 seawater by the marine *Bacillus sp.* strain SG-1. *American Mineralogist* **90**: 1342-1357.

356 Zhang, J.H., Lion, L.W., Nelson, Y.M., Shuler, M.L., Ghiorse, W.C., 2002. Kinetics of Mn(II) oxidation by
357 *Leptothrix discophora* SS1. *Geochim. Cosmochim. Acta* **66**: 773-781.

358

359


360



361

362 **Figure 1.** Mn(II) loss from solution in laboratory sediment incubation experiments. Black
 363 squares (■) – live sediments under air; white squares (□) – live sediments under N₂; black circles
 364 (●) – killed sediments (γ-irradiated) under air; and, white circles (○) – killed sediments under N₂.
 365 For live sediments Mn(II) was repeatedly added (as MnCl₂) at 120, 460 and 600 h, while killed
 366 reactors were never re-spiked with Mn(II). Experiments were conducted with 1.0 g moist
 367 sediment (0.82 g dry) and 50 mL filter sterilized influent site water. Figure from Tan et al.
 368 (2010).


369



371

372 **Figure 2.** Mn(II) loss from solution under variable partial pressures of oxygen (P_{O_2}). White
 373 circles (○) – incubated under 21% P_{O_2} (i.e. air); black squares (■) – incubated under 10% P_{O_2} :
 374 90% N_2 ; and, black triangles (▲) – incubated under 10% P_{O_2} : 99% N_2 . Mn(II) was repeatedly
 375 added (as $MnCl_2$) at 120, 460 and 600 h. Experiments were conducted with 1.0 g moist sediment
 376 (0.82 g dry) and 50 mL filter sterilized influent site water. Corresponding controls using killed
 377 sediments (γ -irradiated) are not shown.

378



380

381 **Figure 3.** Mn(II) loss from solution with and without supplemental organic carbon source.

382 White circles (○) – incubated with 10 mg C/L of glucose; black squares (■) – incubated with no


383 supplemental carbon; and, black triangles (▲) – incubated with 10 mg C/L of

384 carboxymethylcellulose. All incubations maintained under air. Mn(II) was repeatedly added (as

385 MnCl₂) at 120, 460 and 600 h. Experiments were conducted with 1.0 g moist sediment (0.82 g

386 dry) and 50 mL filter sterilized influent site water. Corresponding controls using killed


387 sediments (γ -irradiated) are not shown.



389

390 **Figure 4.** Mn(II) loss from solution in the presence and absence of fungicides. Black squares
 391 (■) – live sediments incubated with no fungicides; white squares (□) – killed sediments (γ -
 392 irradiated) incubated with no fungicides; black circles (●) – live sediments incubated with 0.2
 393 g/L cyclohexamide and 0.2 g/L pentachloronitrobenzene; and, white circles (○) – killed
 394 sediments incubated with 0.2 g/L cyclohexamide and 0.2 g/L pentachloronitrobenzene. All
 395 incubations maintained under air. Mn(II) was repeatedly added (as MnCl_2) at 120, 460 and 600
 396 h. Experiments were conducted with 1.0 g moist sediment (0.82 g dry) and 50 mL filter
 397 sterilized influent site water.

398



399

400 **Figure 5.** Culture-based bacterial enumerations at the end of select sediment incubations.

401 Numbers of culturable heterotrophic bacteria were determined with K media.




Figure 6. Tolerance of fungal isolates to elevated Mn(II) concentrations. Images show two different Mn(II)-oxidizing fungi isolates grown in petri dishes on agar-based media with increasing (0 to 10,000 μ M) concentrations of dissolved Mn(II). Fungal isolate 1 is *Plectosphaerella cucumerina* and fungal isolate 2 is *Microdochium bolleyi*. Isolates were inoculated by “stabbing” the center of the petri dish, allowing the fungal hyphae to grow radially outwards. The brown color is due to the Mn oxide minerals precipitated on the fungi during growth.

Table 1. Aqueous geochemistry of the Fairview Mn-removal system for samples collected in December 2006. Front, middle and rear refer to locations within the limestone treatment bed with the front location nearest the influent. All dissolved concentrations reported in mg/L.

Analyte	Influent	Front	Middle	Rear
Mn	140	137	107	70
Fe	<0.01	<0.01	<0.01	<0.01
Al	5.1	3.0	0.62	0.07
Ca	230	223	248	260
Mg	245	240	228	208
Si	6.5	7.7	5.6	5.7
Na	50	47.5	45	42.5
K	6.6	6.3	6.15	5.65
SO ₄ ²⁻	2010	2010	2010	2010
Cl ⁻	3.65	2.32	2.85	2.57
pH (standard units)	5.43	5.65	6.11	6.34
Alkalinity (mg CaCO ₃ /L)	32.5	25.6	77.5	100
Hot peroxide acidity (mg CaCO ₃ /L)	373	n/a	n/a	n/a
TOC	2.22	2.02	1.96	1.77
TN	0.30	0.18	0.17	0.17
NO ₃ -N	0.18	0.10	0.13	0.13
PO ₄ -P (mg P/L)	<0.1	<0.1	<0.1	<0.1
Dissolved oxygen (mg O ₂ /L)	5.5	7.2	6.1	4.1
Temperature (°C)	10.4	7.1	5.2	4.3